

MINI-SYMPOSIUM

Rate control in the medical management of atrial fibrillation

A J Camm, I Savelieva, G Y H Lip, on behalf of the Guideline Development Group for the NICE clinical guideline for the management of atrial fibrillation

Heart 2007;93:35–38. doi: 10.1136/hrt.2006.099903

Two main strategies are available for management of atrial fibrillation: rate control and rhythm control.

The aims of heart rate control in atrial fibrillation are to minimise symptoms associated with excessive heart rates and to prevent tachycardia-associated cardiomyopathy.¹ Rate control involves the use of negatively chronotropic drugs or electrophysiological/surgical interventions to reduce the rapid ventricular rate often found in patients with atrial fibrillation.² Although the atria continue to fibrillate, this strategy is considered an effective treatment as it can improve symptoms and reduce the risk of associated morbidity. However, the risk of stroke and occurrence of thromboembolic events occurring is reduced by giving antithrombotic drugs.

Rhythm control involves the use of electrical or pharmacological cardioversion or electrophysiological/surgical interventions to convert the arrhythmia associated with atrial fibrillation to normal sinus rhythm. Patients who have been successfully cardioverted are generally given antiarrhythmic drugs for the long term to help prevent the recurrence of atrial fibrillation. Rhythm control strategies also require the appropriate antithrombotic treatment to reduce the risk of stroke and thromboembolism.

WHAT CONSTITUTES ADEQUATE RATE CONTROL?

The optimal heart rate in atrial fibrillation is not known. Current guidelines define adequate rate control in atrial fibrillation as maintenance of the ventricular rate response between 60 and 80 beats/min at rest and between 90 and 115 beats/min during moderate exercise.³ A consensus statement has suggested a target heart rate of <90 at rest and <180 bpm during exercise in patients with atrial fibrillation.⁴ However, no controlled clinical trials have validated these target rates for preventing all-cause cardiovascular morbidity or mortality, and such recommendations may be flawed. Few data exist that define the most robust method for the assessment of rate control.

Adequate rate control may encompass more than the prevention of fast ventricular rates. However, few systematic studies have explored the effect of rate-slowing drugs on chronotropic competence in atrial fibrillation. The effect of rate regularisation is rarely acknowledged in practice, despite evidence that the mere presence of irregular R–R intervals may also contribute to impairment of ventricular haemodynamic function irrespective of heart rate.⁵ Furthermore, atrial contraction contributes between 20% and 40% of the total stroke volume. A marked decrease in cardiac output may occur in atrial fibrillation with the loss of atrial contraction, especially in (mostly elderly) patients with impaired diastolic filling, hypertension and left ventricular hypertrophy.

Although rapid ventricular rates in atrial fibrillation can be detrimental, they should not be too slow either. The heart rate in atrial fibrillation should probably be faster than that in sinus rhythm to maintain an equivalent cardiac output.⁶

Rate-control treatment in atrial fibrillation is based mainly on pharmacological depression of conduction through the atrioventricular node, except in the case of Wolff–Parkinson–White syndrome, where an agent that increases the refractoriness of accessory pathway tissue (eg, flecainide) is needed. Urgent rate control may require the administration of intravenous β -blockers (eg, esmolol) or non-dihydropyridine calcium antagonists. Adequate ventricular rate control at rest does not always translate into effective control during activity, especially with digoxin monotherapy.^{7–8} During exercise, vagal tone (and thus the effect of digoxin) is lost; atrioventricular conduction is further enhanced by the increased sympathetic tone and is different from that in sinus rhythm.⁹ Digoxin is therefore unsuitable for rate-control monotherapy, except in sedentary peoples.

β -Blockers and rate-limiting calcium antagonists are often effective as initial monotherapy for rate control, but a combination of drugs is often necessary to achieve adequate rate control.^{10–12} In clinical practice, β -blockers are commonly used with digoxin for control of exercise-induced increased heart rate, but the evidence for the effectiveness of such combination treatment was less good^{8–13} compared with that available for combination treatment with a rate-limiting calcium antagonist and digoxin.^{8–14–15} Table 1 and fig 1 show the recommendations for rate control.

Although serious adverse effects from rate-control drugs are uncommon, the strategy is not without its limitations and drawbacks. Rate control amounts to negative chronotropic treatment, and may result in slow ventricular rates, sinus bradycardia or heart block (in cases of paroxysmal atrial fibrillation). In some patients, particularly elderly patients, symptomatic bradycardia may require permanent pacing. Of particular importance is that the outcome does not seem to be affected by the achieved heart rates.¹⁶

In patients who fail to respond to medical means of rate control, non-pharmacological measures such as atrioventricular nodal ablation can be considered.¹⁷ Notably, non-pharmacological treatments of rate control were not widely used in the major rate-control versus rhythm-control studies; only 1–2% of

Table 1 Recommendations for rate control

In patients with permanent atrial fibrillation, who need treatment for rate control:

- β -blockers or rate-limiting calcium antagonists should be the preferred initial monotherapy in all patients
- digoxin should only be considered as monotherapy in predominantly sedentary patients

In patients with permanent atrial fibrillation, where monotherapy is inadequate:

- to control the heart rate only during normal activities, β -blockers or rate-limiting calcium antagonists should be given with digoxin
- to control the heart rate during both normal activities and exercise, rate-limiting calcium antagonists should be given with digoxin




Figure 1 Rate-control treatment algorithm for permanent (and some cases of persistent) atrial fibrillation (AF). (1) Patients with permanent AF include those with persistent AF who have been selected for a rate-control treatment strategy. (2) Based on stroke risk stratification algorithm. (3) Target a resting heart rate of <90 bpm (110 bpm for those with recent-onset AF). Target an exercise heart rate of <110 bpm (inactive), 200 minus age (active). (4) Referral for further specialist investigation should be considered especially in those with lone AF or electrocardiogram evidence of an underlying electrophysiological disorder (eg, Wolffe–Parkinson–White syndrome) or where pharmacological treatment has failed.

patients assigned to rate control had ablation of the auriculo-ventricular node, usually in patients with disturbing symptoms poorly controlled by rate-slowing drugs.

RATE CONTROL VERSUS RHYTHM CONTROL

Until recently, there were uncertainties about the most appropriate initial treatment strategy—either rate control or rhythm control—for individual patients.

The Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study¹⁸ enrolled 4060 patients with atrial fibrillation over 6 years. Patients were aged at least 65 years or had another risk factor for stroke or death. Virtually any atrial fibrillation was allowed, including first-onset atrial fibrillation and paroxysmal atrial fibrillation with a documented paroxysm of at least 6 h duration. The mean follow-up was 3.5 years, with no noticeable mortality difference with either a rate-control or a rhythm-control strategy. Notably, non-cardiovascular deaths

(including cancer) accounted for most of the excess deaths in the rhythm-control limb. The rhythm-control strategy was associated with a slightly higher incidence of stroke (7.3% *v* 5.7%), probably owing to more frequent discontinuation of anticoagulant treatment, and greater rates of torsade de pointes (0.8% *v* 0.2%) and hospitalisations (80% *v* 73%). The subgroup analyses suggested a better outcome in younger people <65 years (24%) and in patients with left ventricular dysfunction (23%) who were assigned to rhythm control.

The RAtE Control versus Electrical cardioversion (RACE) trial¹⁹ and two pilot studies, Strategies of Treatment of Atrial Fibrillation (STAF)²⁰ and Pharmacological Intervention in Atrial Fibrillation (PIAF),²¹ focused on patients exclusively with persistent atrial fibrillation and pursued an aggressive rhythm-control strategy by means of serial cardioversions and antiarrhythmic drug treatment. The RACE study reported a non-significant trend towards more end points in the rhythm-control arm (22.6% *v* 17.2%), with patients with hypertension (approximately half the total study population) assigned to rhythm control having a markedly higher event rate compared with their counterparts in the rate-control arm (30.8% *v* 17.3%); this relationship was reversed in participants with normotension (12.5% *v* 17.1%). There was no difference in the occurrence of the combined primary end point between rate-control and rhythm-control arms (5.5% *v* 6.1% per year) in the STAF study of 200 patients at high risk.²⁰ In the PIAF study, the rhythm-control strategy resulted in better exercise performance but did not affect symptoms or quality of life, and was associated with an increased number of hospitalisations for repeat cardioversion and the adverse effects of antiarrhythmic drugs.²¹ The HOw to Treat Chronic Atrial Fibrillation (HOT CAFÉ) study enrolled 205 patients with persistent atrial fibrillation.²² During a mean follow-up of 1.7 years, no difference in the primary end point was seen. However, there was a greater incidence of hospitalisations in the rhythm-control arm (1 per patient) compared with 0.05 per patient in the rate-control arm, but 88% of admissions were for repeat cardioversion. Finally, the Control of Rate versus Rhythm in Rheumatic Atrial Fibrillation Trial (CRRRAFT) in young people (mean age 39 years) who underwent surgery for mitral stenosis²³ reported an advantage of rhythm control with amiodarone over rate control with regard to exercise tolerance and quality of life, although there was no difference in hospitalisation rates, bleeding and thromboembolic events during a follow-up of 1 year.

Overview of the studies

No study found rate control to be inferior to rhythm control or vice versa for the outcome measures of mortality^{18–22} or quality of life.^{24–27} However, the AFFIRM study¹⁸ found mortality to be higher for rhythm control in patients with coronary heart disease, those without heart failure and those >65 years old. The same study did not find rhythm control to be associated with lower mortality in those with heart failure. Rates of hospitalisation, as well as rates of adverse events, were higher among participants treated according to rhythm-control plans.^{18–20–22} The results were generally consistent across studies and considered both older participants with increased risk of stroke^{18–19} and younger participants.^{21–22}

It was thought that many of the stroke and thromboembolic events in the rhythm-control arm were related to the lack of appropriate use of antithrombotic treatment. Rates of anticoagulation were often not controlled between the two treatment arms in many of the studies. For example, in the AFFIRM study,¹⁸ although anticoagulation was mandatory in the rate-control treatment arm, the decision to continue anticoagulation in the rhythm-control arm after 4 weeks of cardioversion was

Table 2 Non-pharmacological approaches to rhythm control

As some patients with persistent atrial fibrillation will satisfy criteria for either an initial rate-control or a rhythm-control strategy (eg, age >65 years but also symptomatic):

- the indications for each option should not be regarded as mutually exclusive and the potential advantages and disadvantages of each strategy should be explained to patients before agreeing on which to adopt
- any comorbidities that might indicate one approach rather than the other should be taken into account
- irrespective of whether a rate-control or a rhythm-control strategy is adopted in patients with persistent atrial fibrillation, the appropriate antithrombotic treatment should be used

A rate-control strategy should be the preferred initial option in patients with the following conditions with persistent atrial fibrillation:

- age >65 years
- coronary artery disease
- contraindications to antiarrhythmic drugs
- unsuitable for cardioversion
- without congestive heart failure

A rhythm-control strategy should be the preferred initial option in patients with the following conditions with persistent atrial fibrillation:

- those who are symptomatic
- younger patients
- presenting for the first time with lone atrial fibrillation
- those with atrial fibrillation secondary to a treated or corrected precipitant
- those with congestive heart failure

Patients unsuitable for cardioversion include those with:

- contraindications to anticoagulation
- structural heart disease (eg, large left atrium >5.5 cm, mitral stenosis) that precludes long-term maintenance of sinus rhythm
- a long duration of atrial fibrillation (usually >12 months)
- multiple failed attempts at cardioversion and/or relapses, even with concomitant use of antiarrhythmic drugs or non-pharmacological approaches
- an ongoing but reversible cause of atrial fibrillation (eg, thyrotoxicosis)

left to the treating doctor. Use of anticoagulation treatment may have had some effect on the outcome of the study, and in a subsequent retrospective analysis,²⁸ the use of anticoagulation treatment was an independent predictor of survival.

Although the AFFIRM investigators²⁹ reported a higher incidence of adverse pulmonary events associated with rhythm control, it was not stated whether this was due to the particular choice of rhythm control drug used rather than the strategy itself.

Also, the RACE, STAF, HOT CAFÉ and PIAF studies—which contributed approximately 25% to the total patient population—included patients with persistent atrial fibrillation, often with previous attempts at cardioversion and maintenance of sinus rhythm, and a fairly high risk of recurrence. In the AFFIRM study, 36% of patients were enrolled after the first episode of atrial fibrillation, many had paroxysmal atrial fibrillation and the duration of atrial fibrillation varied from several hours to several years. The differences in a qualifying episode of atrial fibrillation could account for a markedly higher incidence of long-term maintenance of sinus rhythm in the AFFIRM study (60% at 5 years) compared with RACE (40% at 3 years) and STAF (26% at 2 years). In the HOT CAFÉ study, 63.5% were in sinus rhythm at the end of the study, but the duration of follow-up was only 1 year. In the AFFIRM trial, 40% of the rate-control arm reverted spontaneously to sinus rhythm, and the closely similar primary endpoint results for the rhythm control and rate-control strategies may well be due to a general failure to achieve a clear difference with respect to rhythm and rate status in the two arms of the trials.

These data do not mean that rhythm control is not beneficial, but highlight the limitations of current treatments to achieve

and maintain sinus rhythm. Poor efficacy, proarrhythmia and organ toxicity of antiarrhythmic drugs seem to negate the inherent advantage of sinus rhythm over atrial fibrillation.²⁸ Notably, there was a 1.5-fold excess of non-cardiovascular deaths, primarily due to pulmonary causes or cancer, in the rhythm-control arm of the AFFIRM study, but no difference in cardiovascular deaths.²⁹

It is not uncommon for atrial fibrillation to coexist with congestive heart failure. Rate control has been advocated in patients with chronic congestive heart failure primarily owing to the notion that most antiarrhythmic drugs, except for amiodarone, are associated with increased mortality. However, it is unknown whether rhythm control may benefit patients with recent-onset, asymptomatic atrial fibrillation or those with mild left ventricular dysfunction. Retrospective analyses of large randomised controlled mortality studies have shown that sinus rhythm is a predictor of better survival in patients with New York Heart Association (NYHA) II–IV class heart failure irrespective of treatment assignment.^{30–31} A retrospective analysis of >1000 patients with heart failure showed no difference in 2-year survival between patients in whom restoration and maintenance of sinus rhythm was attempted and those managed with rate control.³² However, patients with heart failure were under-represented in the recent rate-control versus rhythm-control studies. For example, in the AFFIRM trial, 23% had a history of congestive heart failure and only 9% had a history of NYHA functional class ≥II, and there was a non-significant trend towards better survival of these patients when assigned to rhythm control.¹⁸ In the RACE study, approximately half the patients had a history of heart failure and NYHA functional class II,¹⁹ and those treated with rate control had more hospitalisations for heart failure.³³ Conversely, patients treated with rhythm control had more thromboembolic events and were more likely to require a pacemaker. The ongoing Atrial Fibrillation and Congestive Heart Failure trial will examine whether the rhythm-control strategy reduces cardiovascular death in patients with NYHA class II–IV heart failure and an ejection fraction ≤35%.³⁴

Quality of life and cost efficacy

Quality of life analyses showed no major advantage of either a rate-control or a rhythm-control strategy probably because of the enrolment of less symptomatic patients, the inability to achieve stable sinus rhythm or adequate rate control in all patients, and the sizeable dropout and “crossover” rates.^{24–27} All studies reported an overall improvement in quality of life with both treatment strategies. The AFFIRM and RACE studies also conducted formal economic analyses and found that rate control was generally more cost effective than rhythm control.^{35–36} Higher costs associated with the primary rhythm-control strategy were driven by hospitalisations for cardioversion and initiation of antiarrhythmic drug treatment, pacemakers, emergency admissions, and outpatient visits to monitor the outcome and costs of antiarrhythmic drug prescriptions.

The health economic evidence from the National Institute for Health and Clinical Excellence guideline concluded that a rate-control strategy was more cost effective than a rhythm-control strategy. However, it was unclear whether the UK setting and outcome costs of inadequate anticoagulation would affect the results.

Therapeutic implications

The results of the rate versus rhythm trials generally pertain to patients with persistent atrial fibrillation, aged >65 years, who are mildly symptomatic, or to patients with persistent recurrent atrial fibrillation refractory to several drugs, but they cannot be

easily extrapolated to younger patients with paroxysmal or first-onset atrial fibrillation. Highly symptomatic patients, especially if they remain symptomatic despite rate control, must be considered for rhythm control. Because the outcome is very similar for both strategies, provided that anticoagulation continues to be used despite apparent resumption of sinus rhythm, doctors should discuss the pros and cons of both strategies with their patients and engage with the patient in making a decision. Non-pharmacological approaches to rhythm control should be considered when rate control is inappropriate (table 2).

Authors' affiliations

A J Camm, **I Savelieva**, Division of Cardiac & Vascular Sciences, St George's University of London, London, UK

G Y H Lip, University Department of Medicine, City Hospital, Birmingham, UK

Competing interests: None declared.

Correspondence to: A J Camm, British Heart Foundation Professor, St George's University of London, Cranmer Terrace, London SW17 0RE, UK; jcam@sul.ac.uk

Published Online First 8 September 2006

REFERENCES

- Umana E, Solares CA, Alpert MA. Tachycardia-induced cardiomyopathy. *Am J Med* 2003;**114**:51–5.
- Freestone B, Lip GYH. Heart rate control. In: Lip GYH, Godtfredsen J, eds. *Cardiac arrhythmias: a clinical approach*. Edinburgh: Mosby, 2003:275–98.
- Fuster V, Ryden LE, Asinger RW, et al. ACC/AHA/ESC guidelines for the management of patients with atrial fibrillation: executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines and Policy Conferences (Committee to Develop Guidelines for the Management of Patients with Atrial Fibrillation): developed in collaboration with the North American Society of Pacing and Electrophysiology. *J Am Coll Cardiol* 2001;**38**:1231–66.
- Boyd KM, Clark DH, Colthart AB, et al. Final consensus statement of the Royal College of Physicians of Edinburgh Consensus Conference on atrial fibrillation in hospital and general practice, 3–4 September 1998. *Br J Haematol* 1999;**104**:195–6.
- Daoud EG, Weiss R, Bahu M, et al. Effect of an irregular ventricular rhythm on cardiac output. *Am J Cardiol* 1996;**78**:1433–6.
- Rawles JM. What is meant by a "controlled" ventricular rate in atrial fibrillation? *Br Heart J* 1990;**63**:157–61.
- Beasley R, Smith DA, McHaffie DJ. Exercise heart rates at different serum digoxin concentrations in patients with atrial fibrillation. *BMJ (Clin Res Ed)* 1985;**290**:9–11.
- Farshi R, Kistner D, Sarma JS, et al. Ventricular rate control in chronic atrial fibrillation during daily activity and programmed exercise: a crossover open-label study of five drug regimens. *J Am Coll Cardiol* 1999;**33**:304–10.
- Li-Saw-Hee FL, Lip GYH. Digoxin revisited. *QJM* 1998;**91**:259–64.
- Khand AU, Rankin AC, Martin W, et al. Carvedilol alone or in combination with digoxin for the management of atrial fibrillation in patients with heart failure? *J Am Coll Cardiol* 2003;**42**:1944–51.
- Segal JB, McNamara RL, Miller MR, et al. The evidence regarding the drugs used for ventricular rate control. *J Fam Pract* 2000;**49**:47–59.
- Olshansky B, Rosenfeld LE, Warner AL, et al. The Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study: approaches to control rate in atrial fibrillation. *J Am Coll Cardiol* 2004;**43**:1201–8.
- Wong CK, Lau CP, Leung WH, et al. Usefulness of labetalol in chronic atrial fibrillation. *Am J Cardiol* 1990;**66**:1212–5.
- Roth A, Harrison E, Mitani G, et al. Efficacy and safety of medium- and high-dose diltiazem alone and in combination with digoxin for control of heart-rate at rest and during exercise in patients with chronic atrial fibrillation. *Circulation* 1986;**73**:316–24.
- Lewis RV, Laing E, Moreland TA, et al. A comparison of digoxin, diltiazem and their combination in the treatment of atrial fibrillation. *Eur Heart J* 1988;**9**:279–83.
- Cooper HA, Bloomfield DA, Bush DE, et al. Relation between achieved heart rate and outcomes in patients with atrial fibrillation (from the Atrial Fibrillation Follow-up Investigation of Rhythm Management [AFFIRM] Study). *Am J Cardiol* 2004;**93**:1247–53.
- Wood MA, Brown-Mahoney C, Kay GN, et al. Clinical outcomes after ablation and pacing therapy for atrial fibrillation: a meta-analysis. *Circulation* 2000;**101**:1138–44.
- The Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) Investigators. A comparison of rate control and rhythm control in patients with atrial fibrillation. *N Engl J Med* 2002;**347**:1825–33.
- Van Gelder IC, Hagens VE, Bosker HA, et al. A comparison of rate control and rhythm control in patients with recurrent persistent atrial fibrillation. *N Engl J Med* 2002;**347**:1834–40.
- Carlsson J, Miketic S, Windeler J, et al. Randomized trial of rate control versus rhythm control in persistent atrial fibrillation: the Strategies of Treatment in Atrial Fibrillation (STAF) Study. *J Am Coll Cardiol* 2003;**41**:1690–6.
- Hohnloser SH, Kuck KH, Lilienthal J. Rhythm or rate control in atrial fibrillation – Pharmacological Intervention in Atrial Fibrillation (PIAF): a randomised trial. *Lancet* 2000;**356**:1789–94.
- Opolski G, Torbicki A, Kosior DA, et al. Rate control vs rhythm control in patients with nonvalvular persistent atrial fibrillation: the results of the Polish How to Treat Chronic Atrial Fibrillation (HOT CAFE) Study. *Chest* 2004;**126**:476–86.
- Vora A, Karnad D, Goyal V, et al. Control of heart rate versus rhythm in rheumatic atrial fibrillation: a randomized study. *J Cardiovasc Pharmacol Ther* 2004;**9**:65–73.
- Gronefeld GC, Lilienthal J, Kuck KH, et al. Pharmacological-Intervention-in-Atrial-Fibrillation. Impact of rate versus rhythm control on quality of life in patients with persistent atrial fibrillation. Results from a prospective randomized study. *Eur Heart J* 2003;**24**:1430–6.
- Carlsson J, Micus S, Haun S, et al. Rate control and rhythm control improve quality of life in patients with atrial fibrillation: results from STAF (Strategies of Treatment of Atrial Fibrillation) [abstract]. *Circulation* 2001;**104**:II-774.
- Hagens VE, Ranchar AV, Van Sonderen E, et al. Effect of rate or rhythm control on quality of life in persistent atrial fibrillation. Results from the Rate Control Versus Electrical Cardioversion (RACE) Study. *J Am Coll Cardiol* 2004;**43**:241–7.
- Jenkins LS, Brodsky M, Schron E, et al. Quality of life in atrial fibrillation: the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study. *Am Heart J* 2005;**149**:112–20.
- Corley SD, Epstein AE, DiMarco JP, et al. Relationships between sinus rhythm, treatment, and survival in the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) Study. *Circulation* 2004;**109**:1509–13.
- Steinberg JS, Sadaniantz A, Kron J, et al. Analysis of cause-specific mortality in the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study. *Circulation* 2004;**109**:1973–80.
- Dries DL, Exner DV, Gersh BJ, et al. Atrial fibrillation is associated with an increased risk for mortality and heart failure progression in patients with asymptomatic and symptomatic left ventricular systolic dysfunction: a retrospective analysis of the SOLVD trials. Studies of Left Ventricular Dysfunction. *J Am Coll Cardiol* 1998;**32**:695–703.
- Swedberg K, Olsson LG, Charlesworth A, et al. Prognostic relevance of atrial fibrillation in patients with chronic heart failure on long-term treatment with beta-blockers: results from COMET. *Eur Heart J* 2005;**26**:1303–8.
- Al-Khatib SM, Shaw LK, Lee KL, et al. Is rhythm control superior to rate control in patients with atrial fibrillation and congestive heart failure? *Am J Cardiol* 2004;**94**:797–800.
- Hagens VE, Crijns HJ, Van Veldhuisen DJ, et al. Rate control versus rhythm control for patients with persistent atrial fibrillation with mild to moderate heart failure: results from the RACE Control versus Electrical cardioversion (RACE) study. *Am Heart J* 2005;**149**:1106–11.
- AF-CHF Investigators. Rationale and design of a study assessing treatment strategies of atrial fibrillation in patients with heart failure: the Atrial Fibrillation and Congestive Heart Failure (AF-CHF) trial. *Am Heart J* 2002;**144**:597–607.
- Hagens VE, Vermeulen KM, TenVergert EM, et al. Rate control is more cost-effective than rhythm control for patients with persistent atrial fibrillation—results from the RACE Control versus Electrical cardioversion (RACE) study. *Eur Heart J* 2004;**25**:1542–9.
- Marshall DA, Levy AR, Vidaillet H, et al. Cost-effectiveness of rhythm versus rate control in atrial fibrillation. *Ann Intern Med* 2004;**141**:653–61.